

The Hubble Constant from the Fornax Cluster Distance [★]

Tom Richtler¹, Georg Drenkhahn^{2,1}, Matías Gómez^{1,3}, and Wilhelm Seggewiss¹

¹ Sternwarte der Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

² Max-Planck-Institut für Astrophysik, D-85740 Garching, Germany

³ Departamento de Astronomía y Astrofísica, PUC, Santiago de Chile

Abstract. Type Ia supernovae are the best cosmological standard candles available. The intrinsic scatter of their decline-rate- and colour-corrected peak brightnesses in the Hubble diagram is within observational error limits, corresponding to an uncertainty of only $3 \text{ km s}^{-1} \text{ Mpc}^{-1}$ of the Hubble constant. Any additional uncertainty, resulting from peak-brightness calibration, must be kept small by measuring distances to nearby host galaxies most precisely.

A number of different distance determinations of the Fornax cluster of galaxies agree well on a distance modulus of $31.35 \pm 0.04 \text{ mag}$ ($18.6 \pm 0.3 \text{ Mpc}$). This leads to accurate absolute magnitudes of the well-observed Fornax type Ia SNe SN 1980N, SN 1981D, and SN 1992A and finally to a Hubble constant of $H_0 = 72 \pm 6 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

1 The Hubble diagram of type Ia SNe

Supernovae (SNe) of type Ia are known to be excellent standard candles. In order to construct a reliable Hubble diagram it needs more than just to plot the data from the standard SNe catalogue [1] but requires highly uniform spectroscopic and, especially, photometric observations as are now available from the Calán-Tololo survey [2] or the CfA campaign [3].

Fig. 1 shows the Hubble diagram for 29 classified Ia SNe from the Calán-Tololo survey [2]. The intersection Z of the straight line (slope 5!) with the y -axis is related to the Hubble constant H_0 and the absolute magnitude M of the standard candles by the well-known equation:

$$\log H_0 = 0.2 \cdot (M - Z) + 5 \quad (1)$$

For the knowledge of an accurate Hubble constant, one would of course aim at determining the distances to as many SNe as possible.

The quality of the SNe Ia Hubble diagram can be improved further if one corrects the brightness at maximum for the decline rate Δm_{15} of the B lightcurve (e.g. [2]) and the colour ($B_{\text{max}} - V_{\text{max}}$) [4]:

$$B_{\text{corr}} = B_{\text{max}} + b(\Delta m_{15} - 1.1) + R(B_{\text{max}} - V_{\text{max}}) \quad (2)$$

[★] to be published in: Science in the VLT Era and Beyond. ESO VLT Opening Symposium, 1-4 March 1999, Antofagasta, Chile. Springer-Verlag, Berlin, in press




Fig. 1. The Hubble diagram of corrected SN Ia magnitudes in B based on the Calán-Tololo sample [2]. b and R are the correction coefficients for decline rate and intrinsic colour, resp., and Z is the intersection of the straight line with the y -axis. No systematic differences are seen between SNe Ia's in early-type galaxies (circles), late-type galaxies (triangles), and “red” SNe (open symbols).

A new analysis of the Calán-Tololo SNe sample has been made recently by Drenkhahn & Richtler [5,6]. The derived correction terms are quoted in the upper part of Fig. 1. As a result of the correction process the scatter in the Hubble diagram is reduced to 0.14 mag (even smaller in V and I), which is already consistent with the observational errors. Therefore, the intrinsic scatter must be distinctly smaller than 0.1 mag. This fixes the zero-point Z with high precision and the accuracy of H_0 in (1) depends only on the absolute magnitude M_{corr} . Determining M_{corr} within ± 0.1 mag corresponds to measuring H_0 with an uncertainty of only $\pm 3 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 The distance of the Fornax cluster of galaxies

With respect to the accurate calibration of the absolute magnitudes of Ia SNe, the Fornax cluster of galaxies is one of the most suitable sites. It hosts the type Ia SNe SN 1992A (in NGC 1380), SN 1980N, and SN 1981D (both in NGC 1316). In contrast to the Virgo cluster, which is the permanent but precarious target of distance measurements, the Fornax cluster has a compact structure without distinct substructures. Promising attempts to tighten the distance to Fornax have been made [8,10,15], but have been criticised when hunting for a small value of the Hubble constant [7]. The last year has brought

convincing new results for the Fornax distance (for the following discussion refer to Table 1).

Planetary nebula luminosity function (PNLF). Distances to Fornax galaxies using the PNLf have been deduced by Jacoby and co-workers [8,9]. Their zero-point is based on M31 with a distance of 710 kpc (or distance modulus $m - M = 24.26$ mag) and a reddening of $E_{B-V} = 0.11$ mag. Changing to the more recent distance modulus of $m - M = 24.44$ mag [17] but keeping the reddening one arrives at a Fornax modulus of 31.33 ± 0.08 mag.

Globular cluster luminosity function (GCLF). The turn-over magnitude (TOM) of the GCLF of four Fornax galaxies has been derived by Kohle et al. [10] to be $V_{TO} = 23.69 \pm 0.06$ mag (error weighted and skipping NGC 1379). Applying a more recent calibration of the TOM of the Galactic globular cluster system: $M_{V,TO} = -7.61 \pm 0.08$ mag [5,7], one gets a Fornax distance modulus of 31.30 ± 0.13 mag. The GCLF of NGC 1380 has also been analysed using ESO NTT data [11], which result in a modulus of 31.35 ± 0.16 mag. Presently, we are investigating the GCLF of the Fornax galaxy NGC 1316 [12]. The preliminary result of 31.32 ± 0.15 mag agrees well with the other values.

Cepheid distances. Now there is also a Cepheid distance of one Fornax galaxy, NGC 1365, available (HST Key Project [13]). The value of 31.35 ± 0.07 mag agrees very well with the results of the previously mentioned methods.

Surface brightness fluctuations (SBF). Jensen et al. [14] report new infrared SBF measurements for five Fornax galaxies. They find a distance modulus of 31.37 ± 0.12 mag.

Tully-Fisher relation (T-F). Finally, a Tully-Fisher distance from 21 cm data for 18 Fornax galaxies can be derived: when combining published 21 cm line widths ΔV_{20} [15] with dereddened integrated B magnitudes B_{tot}^0 [16] one finds the relation: $B_{tot}^0 = -6.990(\pm 0.426) \log \Delta V_{20} + 29.312(\pm 0.365)$ (N.B.: The galaxy ESO 357-G25 which has an extremely large error in the line width has been skipped from the sample [15, their Table 1].) We then use the absolute magnitudes M_B of all non-Fornax calibrating galaxies from the work of Federspiel et al. [17, their Table 2] and obtain 31.43 ± 0.12 mag.

Table 1 suggests that the distance of the Fornax cluster is now very well determined. The mean value of the distance modulus is 31.35 mag with a scatter of only ± 0.04 mag (linear distance 18.6 ± 0.3 Mpc).

3 The absolute magnitudes of type Ia SNe in Fornax and the Hubble constant

Table 2 lists the parameters of the three Fornax Ia events. The photometry is taken from [2]. For shortness, only the brightness B is given. SN 1992A in NGC 1380 is one of the best ever observed SNe. After correction for decline rate and colour and with the new distance modulus of the Fornax cluster the

Table 1. Distance modulus determinations of the Fornax cluster

Galaxy NGC	$m - M$ mag	σ mag	Method ^a	Reference
1316, 1399, 1404	31.33	0.08	PNLF	[8], [9]
1374, 1399, 1427	31.30	0.11	GCLF	[10]
1380	31.35	0.16	GCLF	[11]
1316	31.32	0.15	GCLF	[12]
1365	31.35	0.07	Ceph	[13]
1339, 1344, 1379, 1399, 1404	31.37	0.12	SBF	[14]
18 galaxies	31.43	0.12	T-F	[15], [16], [17]
mean value	31.35	0.04		

- ^a PNLf: planetary nebulae luminosity function
GCLF: globular cluster luminosity function
Ceph: Cepheid distance; HST Key Project
SBF: surface brightness fluctuations
T-F: Tully-Fisher relation

absolute magnitudes M_{corr}^B have been calculated. Eq. (1) and the corresponding data in the B , V , and I bands for the listed SNe yield a Hubble constant of $H_0 = 72 \pm 6 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

Table 2. Type Ia SNe in the Fornax cluster of galaxies

SN	Host Galaxy	B_{max}	σ	Δm_{15}	$B_{\text{max}} - V_{\text{max}}$	B_{corr}	M_{corr}^B
SN 1992A	NGC 1380	12.57	0.03	1.47	0.02	12.36	-18.99
SN 1980N	NGC 1316	12.49	0.03	1.28	0.05	12.33	-19.02
SN 1981D	NGC 1316	12.66	0.06	1.20	0.24	12.25	-19.10

Parameters of SN 1981D from own light-curve fit

References

- Barbon, R., Cappellaro, E., Turatto, M. (1989) A&AS **81**, 421, <http://athena.pd.astro.it/~supern/>
- Hamuy, M., Phillips, M. M., Suntzeff, N. B., Schommer, R. A., Maza, J., Avilés, R. (1996) AJ **112**, 2398
- Riess, A. G., Kirshner, R. P., Schmidt, B. P. et mult. al. (1999) AJ **117**, 707
- Tripp, R. (1998) A&A **331**, 815
- Drenkhahn, G., Richtler, T. (1999) A&A submitted
- Richtler, T., Drenkhahn, G. (1999) in: Kundt, W., van de Bruck, C. (Eds.) Cosmology and Astrophysics: A Collection of Critical Thoughts. Lecture Notes in Physics, Springer, Berlin, in press

7. Tammann, G. A., Sandage, A. (1999) in: Egret, D., Heck, A. (Eds.) Harmonizing Cosmic Distance Scales in a Post-Hipparcos Era. Obs. Strasbourg, in press
8. McMillan, R., Ciardullo, R., Jacoby, G. H. (1993) *ApJ* **416**, 62
9. Jacoby, G. H. (1997) in: Livio, M., Donahue, M., Panagia, N. (Eds.) The Extragalactic Distance Scale. University Press, Cambridge, p. 197
10. Kohle, S., Kissler-Patig, M., Hilker, M., Richtler, T., Infante, L., Quintana, H. (1996) *A&A* **309**, L39
11. Della Valle, M., Kissler-Patig, M., Danziger, J., Storm, J. (1998) *MNRAS* **299**, 267
12. Gómez, M., Richtler, T., Infante, L. (1999) in preparation
13. Madore, B. F., Freedman, W. L., Silbermann, N. et al. (1998) *Nature* **395**, 47
14. Jensen, J.B., Tonry, J.L., Luppino, G.A. (1998) *ApJ* **505**, 111
15. Bureau, M., Mould, J. R., Staveley-Smith, L. (1996) *ApJ* **463**, 60
16. de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Buta, R. J., Paturel, G., Fouqué, G. (1991) *Third Ref. Cat. of Bright Galaxies*. Springer, New York
17. Federspiel, M., Tammann, G. A., Sandage, A. (1998) *ApJ* **495**, 115